arXiv:1304.4422v1 [math.AT] 16 Apr 2013

COMPUTING KRICHEVER GENUS

MALKHAZ BAKURADZE

Abstract. We prove that on the rational complex bordism ring the Krichever-Höhn genus ϕ_{KH} is the composition $\psi \circ \kappa^{-1}$. Here the genus ψ corresponds to the formal group law whose invariant differential $\omega(t)$ equals $\sqrt{1+p_1t+p_2t^2+p_2t^3+p_4t^4}$ and κ classifies the formal group law strictly isomorphic to the universal formal group law under strict isomorphism $x\mathbf{CP}(x)$. We construct certain elements A_{ij} in the Lazard ring and give an alternative definition of the universal Krichever formal group law. This implies that the corresponding coefficient ring is the quotient of the Lazard ring by the ideal generated by all A_{ij}, $i, j \geq 3$.

The general four variable complex elliptic genus ϕ_{KH} on $MU_* \otimes \mathbb{Q}$ is defined [6],[7] to be determined by the following property: if one denotes by $f_{Kr}(x)$ the exponent of the Krichever [5] universal formal group law $\mathcal{F}_{Kr}$, then the series

$$h(x) := \frac{f'_{Kr}(x)}{f_{Kr}(x)}$$

satisfies the differential equation

$$(h'(x))^2 = S(h(x)), \tag{0.1}$$

where $S(x) = x^4 + p_1x^3 + p_2x^2 + p_2x + p_4$, the generic monic polynomial of degree 4 with formal parameters p_i of weights $|p_i| = 2i$.

To generalize naturally the Oshanin's elliptic genus from $\Omega_*^{SO} \otimes \mathbb{Q}$ to $\mathbb{Q}[\mu, \epsilon]$ [8], new elliptic genus ψ is defined in [9] to be the genus $\psi : MU_* \otimes \mathbb{Q} \to \mathbb{Q}[p_1, p_2, p_3, p_4]$ whose logarithm equals

$$\int_0^x \frac{dt}{\omega(t)}, \quad \omega(t) = \sqrt{1+p_1t+p_2t^2+p_2t^3+p_4t^4}$$

and p_i are again formal parameters $|p_i| = 2i$.

Clearly to calculate the values of ψ on $\mathbf{CP}_i$, the generators of the rational complex bordism ring $MU_* \otimes \mathbb{Q} = \mathbb{Q}[\mathbf{CP}_1, \mathbf{CP}_2, \dots]$ we need only the Taylor expansion of $(1+y)^{-1/2}$ as by above definition

$$(1+p_1x+p_2x^2+p_2x^3+p_4x^4)^{-1/2} = log'_\psi = \sum_{i\geq 1} \psi(\mathbf{CP}_i)x^i.$$

It is natural to ask if there exists similar elementary way, different from formulas in [3], for calculation of ϕ_{KH}. Our result below is the formula of Theorem 0.1.

Let $\mathbf{CP}(x) = 1 + \sum_{i\geq 1} \mathbf{CP}_i x^i$ and let $\nu(x) := x\mathbf{CP}(x)$ be the strict isomorphism $F \to \tilde{F}$. So that

$$\nu(F(x,y)) = \tilde{F}(\nu(x), \nu(y)),$$

where F is the universal formal group law. Let $log(x)$ and $mog(x)$ be the logarithm series of F and $\tilde{F}$ respectively then

$$mog(x) = log(\nu^{-1}(x)).$$

Then let

$$\kappa : MU_* \otimes \mathbb{Q} \to MU_* \otimes \mathbb{Q}$$

be the classifying map of $\tilde{F}$. Then $\kappa(log(x)) = mog(x)$ therefore $\kappa(\mathbf{CP}_i)$ is determined by equating coefficients at x^i in

---

The author was supported by Shota Rustaveli NSF, DI/16/5-103/12 and by Volkswagen Foundation, Ref.: I/84 328 .

$$\sum_{i\geq 1}\frac{\kappa(\mathbf{CP}_i)}{i+1}x^{i+1}=\sum_{i\geq 1}\frac{\mathbf{CP}_i}{i+1}(\nu^{-1}(x))^{i+1}. \tag{0.2}$$

For instance

$\kappa(\mathbf{CP}_1)=-\mathbf{CP}_1$;
$\kappa(\mathbf{CP}_2)=3\mathbf{CP}_1^2-2\mathbf{CP}_2$;
$\kappa(\mathbf{CP}_3)=-10\mathbf{CP}_1^3+12\mathbf{CP}_1\mathbf{CP}_2-3\mathbf{CP}_3$;
$\kappa(\mathbf{CP}_4)=35\mathbf{CP}_1^4-60\mathbf{CP}_1^2\mathbf{CP}_2+20\mathbf{CP}_1\mathbf{CP}_3+10\mathbf{CP}_2^2-4\mathbf{CP}_4$.

Theorem 0.1. $\phi_{KH}=\psi\circ\kappa^{-1}$.

We need the following

Lemma 0.2. *Let F be the universal formal group law and $\tilde{F}$ be its isomorphic under the strict isomorphism $\nu(x)=x\mathbf{CP}(x)$ as above. Let exp be the exponent of F. The series $\frac{1}{h(x)}=exp(x)/exp'(x)$ is reversible, and let $j(x)$ be its reverse, then $j(x)=mog(x)$, the logarithm series of $\tilde{F}$.*

Proof. Note that

$$j^{-1}(log(x))=\frac{exp(log(x))}{exp'(log(x))}=x\mathbf{CP}(x)$$

hence by definition of $\nu(x)$ one has

$$j^{-1}(mog(x))=j^{-1}(log(\nu^{-1}(x)))=\nu^{-1}(x)\mathbf{CP}(\nu^{-1}(x))=x$$

as $x\mathbf{CP}(x)=\nu(x)$. □

The following lemma completes the proof of Theorem 0.1

Lemma 0.3. *Let $\tilde{\omega}=\frac{1}{mog'(x)}$ be the invariant differential form of the formal group law $\tilde{F}$ above. Then the condition of Krichever-Höhn (0.1) is satisfied if and only if $\tilde{\omega}(x)^2$ is a polynomial of degree 4.*

Proof. Since the series $j(x)$ is reversible, the condition (0.1) is equivalent to

$$(h'(j(x)))^2=S(h(j(x))).$$

But $h(j(x))=1/x$, hence

$$(h\circ j)'(x)=h'(j(x))j'(x)=-1/x^2,$$

so that

$$h'(j(x))=-\tilde{\omega}(x)/x^2.$$

It follows that the condition (0.1) is equivalent to

$$\tilde{\omega}(x)^2=x^4S(1/x)$$

where on the right hand side one clearly has a degree four polynomial. □

Now let us turn to universal Krichever formal group law and give its alternative equivalent definition announced in [2].

Let $F(x,y)$ be the formal group law of geometric cobordism [11]. Following Quillen [12] we will identify it with the universal Lazard formal group law. Let $\omega(x)=\frac{\partial F(x,y)}{\partial y}(x,0)$.

In [4] V. M. Buchsteber has given the analytical solution of a functional equation for the exponent of the formal group law of the form

$$\mathcal{F}_B(x,y)=\sum\alpha_{ij}x^iy^j=\frac{A(y)x^2-A(x)y^2}{B(y)x-B(x)y}. \tag{0.3}$$

Lemma 0.4. *Let $\mathcal{F}$ be the formal group law of the form $\mathcal{F}_B$, with $B'(0) = A'(0)$. Then $B(x)$ coincides with restricted $\omega(x)$, that is $B(x)$ is the image of $\omega(x)$ under the ring homomorphism of the Lazard ring classifying the formal group law $\mathcal{F}$.*

For this note that if our series A and B have form

$$A(t) = A_0 + A_1 t + A_2 t^2 + O(t^3)$$

and

$$B(t) = B_0 + B_1 t + B_2 t^2 + O(t^3),$$

then

$$\mathcal{F}(x, y) = \frac{A_0}{B_0}(x + y) + O(xy).$$

So if $\mathcal{F}(x, y)$ is a formal group law we must have $A_0 = B_0$, and after dividing the numerator and denominator appropriately we may assume that $A_0 = B_0 = 1$. We then furthermore calculate

$$\mathcal{F}(x, y) = x + y + A_1 xy + \sum_{i=2}^{\infty} B_i(x^i y + xy^i) + O(x^2 y^2).$$

We thus have

$$\alpha_{1i} = B_i,\ i \geq 2,\ \alpha_{11} = A_1 = A'(0) = B'(0)$$

and

$$1 + \sum_{i \geqslant 1} \alpha_{1i} t^i$$

is restricted $(1 + \sum_{i \geqslant 1} [\mathbf{CP}^i] t^i)^{-1} = \omega(t)$. □

Let us now present a minor modification of the analysis of $\mathcal{F}_B$ performed in [10] and as Lemma 0.4 suggests to introduce

$$A(x, y) = \sum A_{ij} x^i y^j = F(x, y)(x\omega(y) - y\omega(x)). \tag{0.4}$$

We define the universal Nadiradze formal group law $\mathcal{F}_N$ by the obvious classifying map of the Lazard ring to its quotient ring by the ideal generated by all A_{ij} with $i, j \geq 3$.

Proposition 0.5. *Let L be the Lazard ring. Then*

i) $\omega'(x) - \omega'(0) = 2x\hat{\omega}(x)$ in $L[[x]]$, where $\hat{\omega}(x) = \sum_{i \geqslant 1} \omega_i x^{i-1}$.

ii) One has in $L[[x, y]]/(xy)^3$

$$A(x, y) = (x\omega(y) + y\omega(x) - \omega'(0)xy)\,(x\omega(y) - y\omega(x)) + (\omega(x)\hat{\omega}(x) - \omega(y)\hat{\omega}(y))\, x^2 y^2.$$

Proof. Let f and g be the exponent and logarithm of F respectively. Hence $F(x, y) = f(g(x)+g(y))$ and

$$f'(x) = 1/g'(f(x)) = \omega(f(x)),\ \ f'(g(x)) = \omega(f(g(x)) = \omega(x). \tag{0.5}$$

Let $\omega(x) = 1 + b_1 x + b_2 x^2 + \cdots$. Then as $g''(0) = -f''(0) = -\omega'(0) = -b_1$ we have

$$\frac{\partial^2 F}{\partial y^2}(x, 0) = f^{''}(g(x)) + f'(g(x))g''(0) = \omega'(x)\omega(x) - \omega'(0)\omega(x). \tag{0.6}$$

This implies i) as the left hand side of (0.6) has factor 2, and $\omega(x)$ is invertible. ii) Because of antisymmetry we have modulo $(xy)^3$

$$A(x, y) = A(y)x^2 - A(x)y^2 = \sum (A_{i2} x^2 y^i - A_{i2} x^i y^2).$$

We want to calculate $-\sum A_{i2} x^i$ in terms of $\omega(x)$.

Applying $\frac{\partial^2}{\partial y^2}(x, 0)$ to (0.4) and taking into account (0.5) and (0.6) one obtains

$$-2\sum A_{i2} x^i = x\omega(x)\omega'(x) - x\omega'(0)\omega(x) + 2x\omega'(0)\omega(x) - 2\omega^2(x) + 2b_2 x^2.$$

As the coefficients are in the Lazard ring this implies

$$-\sum A_{i2}x^i = x\omega(x)\frac{\omega'(x)-\omega'(0)}{2} + x\omega'(0)\omega(x) - \omega^2(x) + b_2x^2.$$

Therefore

$\sum(A_{i2}x^2y^i - A_{i2}x^iy^2) =$
$(x\omega(y)+y\omega(x))(x\omega(y)-y\omega(x)) - \omega'(0)xy(x\omega(y)-y\omega(x)) + \omega(x)\hat{\omega}(x)x^2y^2 - \omega(y)\hat{\omega}(y)x^2y^2.$ □

In order to compute the Krichever genus on the coefficients of the formal group law of geometric cobordism in [5] the universal Krichever formal group law $\mathcal{F}_{Kr}$ is defined as

$$\mathcal{F}_{Kr}(x,y) = xb(y) + yb(x) - b'(0)xy + \frac{b(x)\beta(x) - b(y)\beta(y)}{xb(y) - yb(x)}x^2y^2, \tag{0.7}$$

where $\beta(x) = \frac{b'(x)-b'(0)}{2x}$. Also it is proved in [5] that

$$b(x) = \frac{\partial \mathcal{F}_{Kr}}{\partial y}(x,0).$$

Lemma 0.4 and Proposition 0.5 ii) imply that $\mathcal{F}_{Kr}$ can be alternatively defined by the classifying map of $\mathcal{F}_N$ if relabel the restriction of the invariant form $\omega(x)$ as $b(x)$. Thus $\mathcal{F}_N = \mathcal{F}_{Kr}$.

Following [4] in [3] the authors consider the following formal group law corresponding to Krichever genus

$$\mathcal{F}_b(u_1,u_2) = u_1c(u_2) + u_2c(u_1) - au_1u_2 - \frac{d(u_1)-d(u_2)}{u_1c(u_2)-u_2c(u_1)}u_1^2u_2^2.$$

It follows from Lemma 0.4 and Proposition 0.5 ii) that (if specify $c'(0) = a$) $\mathcal{F}_b(x,y)$ coincides with (0.7), that is $c(x) = b(x)$ and $d(x) = -b(x)\beta(x)$.

Theorem 0.6. *$\mathcal{F}_N = \mathcal{F}_{Kr} = \mathcal{F}_b$ and the coefficient ring is the the quotient of the Lazard ring by the ideal generated by all A_{ij} with $i,j \geq 3$.*

In [1] we calculated on MAPLE the coefficient ring of Nadiradze formal group law $\mathcal{F}_N$ up to dimension 26. Namely there is a set of polynomial generators $z_1, z_2, \ldots$ of the Lazard ring for which the low degree defining relations are

$$5z_5 = z_2z_3 + 2z_1z_4,\ 2z_6 = 0,\ z_1z_6 = 0,\ z_3z_6 = 0,\ z_{10} = 0,\ z_5z_6 = 0,\ z_{12} = 0,$$

and $7z_7$, $2z_8$, $3z_9$, $11z_{11}$, and $13z_{13}$ are decomposable. For the space reason we omit here these long decompositions. Note that our calculations agree with the results in [5] on the structure of the coefficient ring of $\mathcal{F}_{Kr}$ obtained in terms of associativity equation. The new information here concerning the Krichever group, and hence the Krichever genus, is that as the relations $z_{10} = 0$, $z_{12} = 0$ imply in dimensions 20 and 24 there is no indecomposable elements. The question arises in which dimensions any element is multiplicatively decomposable.

References

[1] M. Bakuradze and M. Jibladze, *On the coeficient ring of rational formal group law*, Proc, A. Razmadze Math Inst. 159, (2012) 1-9.

[2] M. Bakuradze, *Formal group laws by Buchstaber, Krichever and Nadiradze coincide*, Russian Math. Surv., to appear

[3] Victor Buchstaber, Taras Panov, and Nigel Ray, *Toric genera*, Intern. Math. Research Notices, Vol 2010 No. 16, 3207-3262.

[4] V. M. Buchstaber, *Functional equations associated with addition theorems for elliptic functions and two-valued algebraic groups,* Uspekhi Mat. Nauk, Vol 45, Issue 3(273), (1990)185186.

[5] V. M. Buchstaber, E.Yu. Bunkova, *Krichever Formal groups*, Funct. Anal. and appl. 45:2, (2011)23-44 .

[6] Gerald Höhn, emphKomplexe elliptische Geschlechter und S^1-äquivariante Kobordismustheorie, Bonn und Vallendar, August 1991. Available as `arXiv:math/0405232`

[7] I. Krichever, *Generalized elliptic genera and Baker-Akhiezer functions,* Math. Notes 47 (1990), 132-142.

[8] S. Ochanine, *Sur les genres multiplicatifs definis par des integrales elliptiques,* Topology 26 No. 2 (1987), 143-151.

[9] Stefan Schreieder, *Dualization invariance and a new complex elliptic genus*, arXiv:1109.5394v3 [math.AT] 14 Sep 2012

[10] R. Nadiradze, *Formal group and cohomology theories*, Dissertation for the Doct. of Sci. Degree, Tbilisi 1995.

[11] S. P. Novikov, *The methods of algebraic topology from the viewpoint of cobordism theory,* Math. USSR-Izv. vol. 1, issue 4 (1967)827-913

[12] D. Quillen, *On the formal group laws of unoriented and complex cobordism theory*, Bull. Amer. Math. Soc. 75 (1969)1293-1298

Faculty of exact and natural sciences, Iv. Javakhishvili Tbilisi State University, Georgia
E-mail address: `malkhaz.bakuradze@tsu.ge`